

# *Orchis purpurea*

## Habitus and growth type

Plant height [m]: **0.45**

Life span: **Perennial**

Life form: **Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **29.16**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.5**

Temperature indicator value: **6.9**

Moisture indicator value: **4.3**

Reaction indicator value: **8**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.22**

Disturbance frequency (herb layer): **1.44**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.18**

Grazing pressure: **0.24**

Soil disturbance: **0.11**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [\*\*CB \(PUB\) \*Quercetea pubescentis\*\*\*](#), [\*\*DA \(FES\) \*Festuco-Brometea\*\*\*](#)

### **Broad habitat**

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

## **Distribution**

Continentality: **5**

Continentality amplitude: **3**