

# *Orlaya daucoides*

## Habitus and growth type

Plant height [m]: **0.25**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **16.55**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **7.5**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **5.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.5**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.16**

Grazing pressure: **0.3**

Soil disturbance: **0.27**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) \*Chenopodietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#)

Constant species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**