

# *Ornithogalum narbonense*

## Habitus and growth type

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **3.22**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **8.5**

Moisture indicator value: **4.3**

Reaction indicator value: **7.3**

Nutrient indicator value: **5.9**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.08**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.8**

Disturbance severity (herb layer): **0.73**

Mowing frequency: **0.43**

Grazing pressure: **0.25**

Soil disturbance: **0.42**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FF \(LYG\) Lygeo sparti-Stipetea tenacissimae](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**