

# *Ornithopus pinnatus*

## Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **9**

Reaction indicator value: **4.3**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **1.52**

Disturbance frequency (herb layer): **2.05**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.5**

Grazing pressure: **0.28**

Soil disturbance: **0.28**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

Constant species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**