Orobanche alsatica

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**
Moisture indicator value: **3.7**
Reaction indicator value: **7.2**
Nutrient indicator value: **2.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.02**
Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.52**
Grazing pressure: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz