Orobanche latisquama

Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **8.3**
Moisture indicator value: **2.3**
Reaction indicator value: **7**
Nutrient indicator value: **2.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.51**
Disturbance frequency (herb layer): **1.72**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0**
Grazing pressure: **0.39**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz