Ostrya carpinifolia

Habitus and growth type

Plant height [m]: **10.19**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **25.5**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Anemochory**
Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **7.9**
Moisture indicator value: **4.2**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.03**
Disturbance frequency (herb layer): **1.3**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0.01**
Grazing pressure: **0.21**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1G Alnus cordata forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**
Constant species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1G Alnus cordata forest**, **T33 Mediterranean mountain Abies forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**
Dominant species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest**, **T3C Taxus baccata forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz