

# *Osyris lanceolata*

## Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **2.84**

## Fruit, seed and dispersal

Seed mass [mg]:

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Disturbance indicator values

Disturbance frequency: **0.4**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.04**

Grazing pressure: **0.31**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#), [GB \(OLE\)](#)

[Oleo cerasiformis-Rhamnetea crenulatae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U3C Macaronesian inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**