Osyris lanceolata

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm^2/mg]: **2.84**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.4**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.61**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.04**
Grazing pressure: **0.31**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis*, GB (OLE) *Oleo cerasiformis-Rhamnetea crenulatae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U3C Macaronesian inland cliff**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz