

# *Oxalis dillenii*

## Habitus and growth type

Plant height [m]: **0.16**

Life span: **Annual, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## Flower

Flowering period: **January-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.19**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.2**

Moisture indicator value: **5**

Reaction indicator value: **6**

Nutrient indicator value: **5.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.88**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.59**

Mowing frequency: **0.34**

Grazing pressure: **0.28**

Soil disturbance: **0.46**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

## **Distribution**

Continentality amplitude: **5**