

Pallenis spinosa

Habitus and growth type

Plant height [m]: **0.29**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Spinescence: **spinescent**

Leaf

Specific leaf area [mm²/mg]: **15.36**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.8**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8.8**

Moisture indicator value: **2.8**

Reaction indicator value: **7.7**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.86**

Disturbance frequency (herb layer): **1.64**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.62**

Mowing frequency: **0.38**

Grazing pressure: **0.34**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FF \(LYG\) *Lygeo sparti-Stipetea tenacissimae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T26 *Phoenix canariensis* vegetation](#)

Constant species of EUNIS habitats: [R1E Mediterranean tall perennial dry grassland](#), [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#), [T26 *Phoenix canariensis* vegetation](#), [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**