

Pancratium maritimum

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **45.7**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **10.5**

Moisture indicator value: **3.5**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **1.5**

Disturbance indicator values

Disturbance frequency: **1.5**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.01**

Grazing pressure: **0.14**

Soil disturbance: **0.61**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**JD \(AMM\) *Ammophiletea arundinaceae***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**N14 Mediterranean, Macaronesian and Black Sea shifting coastal dune**](#), [**N16 Mediterranean and Macaronesian coastal dune grassland \(grey dune\)**](#), [**N22 Mediterranean and Black Sea coastal shingle beach**](#)

Constant species of EUNIS habitats: [**N14 Mediterranean, Macaronesian and Black Sea shifting coastal dune**](#), [**N16 Mediterranean and Macaronesian coastal dune grassland \(grey dune\)**](#), [**N22 Mediterranean and Black Sea coastal shingle beach**](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle**