

# *Papaver albiflorum*

## Habitus and growth type

Plant height [m]: **0.5**

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **6.3**

Moisture indicator value: **4.6**

Reaction indicator value: **7**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.89**

Disturbance frequency (herb layer): **2.21**

Disturbance severity: **0.85**

Disturbance severity (herb layer): **0.84**

Mowing frequency: **0.22**

Grazing pressure: **0.12**

Soil disturbance: **0.74**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) Koelerio-Coryneporetea  
canescentis](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**