Papaver rhoeas

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **23.89**

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.25**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships
Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**
Ellenberg-type indicator values
Light indicator value: **7.1**
Moisture indicator value: **4.4**
Reaction indicator value: **7**
Nutrient indicator value: **5.9**
Salinity indicator value: **0.1**
Disturbance indicator values
Disturbance frequency: **1.75**

Disturbance frequency (herb layer): **2**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.52**

Grazing pressure: **0.15**

Soil disturbance: **0.74**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QA (PAR) *Papaveretea rhoeadis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**

Constant species of EUNIS habitats: **V11 Intensive unmixed crops**, **V12 Mixed crops of market gardens and horticulture**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**, **V32 Mediterranean subnitrophilous annual grassland**

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality amplitude: **5**