Parentucellia latifolia

Habitus and growth type

Plant height [m]: **0.14**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.8**
Moisture indicator value: **3**
Reaction indicator value: **5.7**
Nutrient indicator value: **3.7**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.34**
Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.62**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.31**
Grazing pressure: **0.43**
Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FH (BUL) *Poetea bulbosae*, FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1D Mediterranean closely grazed dry grassland, S34 Balkan-Anatolian submontane genistoid scrub**

Constant species of EUNIS habitats: **R1D Mediterranean closely grazed dry grassland, S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz