

Pedicularis lapponica

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **16.8**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.24**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0**

Grazing pressure: **0.32**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**AC \(LOI\) Loiseleurio procumbentis-Vaccinietea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S11 Shrub tundra](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [S11 Shrub tundra](#), [S12 Moss and lichen tundra](#), [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Heathland, Forest, Sparsely vegetated (incl. rock and scree)**