

# ***Pedicularis sceptrum-carolinum***

## **Habitus and growth type**

Plant height [m]: **0.58**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **15.58**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.38**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8**

Temperature indicator value: **4.9**

Moisture indicator value: **7.8**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **1.21**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.18**

Mowing frequency: **0.23**

Grazing pressure: **0.11**

Soil disturbance: **0.1**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) \*Scheuchzerio palustris\*-\*Caricetea fuscae\*](#)

Broad habitat

Occurrence in broad habitats: **Mire**

## Distribution

Continentality: **8**

Continentality amplitude: **4**