

# ***Pedicularis verticillata***

## **Habitus and growth type**

Plant height [m]: **0.17**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **April-August**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.8**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8.2**

Temperature indicator value: **2.8**

Moisture indicator value: **6.6**

Reaction indicator value: **7.4**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **1.19**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.04**

Grazing pressure: **0.24**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline)**

## Distribution

Continentality: **6**

Continentality amplitude: **6**