Periploca graeca

Habitus and growth type

Plant height [m]: **2.75**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **11.41**
Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.5**
Moisture indicator value: **7**
Reaction indicator value: **6.5**
Nutrient indicator value: **5.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.33**
Disturbance frequency (herb layer): **1.48**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.32**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.04**
Grazing pressure: **0.19**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FD (NER) *Nerio-Tamaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1J Mediterranean and Black Sea moist and wet dune slack**
Constant species of EUNIS habitats: **N1J Mediterranean and Black Sea moist and wet dune slack**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz