Petrosimonia oppositifolia

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5**
Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **2.17**
Disturbance frequency (herb layer): **2.17**
Disturbance severity: **0.4**
Disturbance severity (herb layer): **0.4**
Mowing frequency: **0.03**
Grazing pressure: **0.49**
Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***, **EA (LER) *Artemisietea lerchianae***, **EC (KAL) *Kalidietea foliati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R64 Semi-desert salt pan**, **S67 Aralo-Caspian semi-desert**, **S94 Semi-desert riparian scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Constant species of EUNIS habitats: **R64 Semi-desert salt pan**, **S67 Aralo-Caspian semi-desert**, **S94 Semi-desert riparian scrub**

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz