

# *Peucedanum altissimum*

## Habitus and growth type

Plant height [m]: **2.1**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5**

Temperature indicator value: **6**

Moisture indicator value: **7**

Reaction indicator value: **6.5**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.26**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.08**

Grazing pressure: **0.22**

Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#), [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**