

# *Phelipanche purpurea*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **6.6**

Moisture indicator value: **3.7**

Reaction indicator value: **7.2**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.34**

Grazing pressure: **0.28**

Soil disturbance: **0.21**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [\*\*DA \(FES\) Festuco-Brometea\*\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

## **Distribution**

Continentality: **6**

Continentality amplitude: **6**