Phleum ambiguum

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **6.4**
Moisture indicator value: **2.7**
Reaction indicator value: **6.5**
Nutrient indicator value: **2.2**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.07**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.43**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.17**
Grazing pressure: **0.36**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**, **S74 Central Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Peninsula, S74 Central Mediterranean mountain hedgehog-heath

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz