

# *Phlomis lanata*

## Habitus and growth type

Plant height [m]: **0.75**

Life span: **Perennial**

Life form: **Phanerophyte, Chamaephyte, Shrub**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **4.6**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.25**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.02**

Grazing pressure: **0.32**

Soil disturbance: **0.18**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

Constant species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**