

Phyteuma globulariifolium

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **1.81**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.05**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **1.3**

Moisture indicator value: **4**

Reaction indicator value: **2.5**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.43**

Disturbance frequency (herb layer): **1.51**
Disturbance severity: **0.26**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0**
Grazing pressure: **0.23**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CT \(TR1\) *Juncetea trifidi*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Constant species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**