

# *Pimpinella nigra*

## Habitus and growth type

Plant height [m]: **0.43**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **6**

Moisture indicator value: **3**

Reaction indicator value: **6.7**

Nutrient indicator value: **2.3**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.98**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.65**

Grazing pressure: **0.28**

Soil disturbance: **0.14**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **6**

Continentality amplitude: **3**