Pistacia lentiscus

Habitus and growth type

Plant height [m]: **3.75**
Life span: **Perennial**
Life form: **Phanerophyte, Tree, Shrub**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **5.44**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **27.04**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **9.5**
Moisture indicator value: **3**
Reaction indicator value: **6.3**
Nutrient indicator value: **4.5**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.23**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.01**
Grazing pressure: **0.28**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1B Mediterranean and Black Sea coastal dune scrub**, **N1G Mediterranean coniferous coastal dune forest**, **S51 Mediterranean maquis and arborescent matorral**, **S54 Thermomediterranean arid scrub**, **S62 Western acidophilous garrigue**, **S71 Western Mediterranean spiny heath**, **T24 Olea europaea-Ceratonia siliqua forest**, **T25 Phoenix theophrasti vegetation**, **T3A Mediterranean lowland to submontane Pinus forest**
Constant species of EUNIS habitats: **N1B Mediterranean and Black Sea coastal dune scrub**, **N1G Mediterranean coniferous coastal dune forest**, **S51 Mediterranean maquis and arborescent matorral**, **S53 Spartium junceum scrub**, **S54 Thermomediterranean arid scrub**, **S62 Western acidophilous garrigue**, **S63 Eastern garrigue**, **S71 Western Mediterranean spiny heath**, **S72 Eastern Mediterranean spiny heath (phrygana)**, **S93 Mediterranean riparian scrub**, **T1A Mediterranean thermophilous deciduous forest**, **T21 Mediterranean evergreen Quercus forest**, **T24 Olea europaea-Ceratonia siliqua forest**, **T25 Phoenix theophrasti vegetation**, **T3A Mediterranean lowland to submontane Pinus forest**
Dominant species of EUNIS habitats: **N1B Mediterranean and Black Sea coastal dune scrub**, **N1G Mediterranean coniferous coastal dune forest**, **S51 Mediterranean maquis and arborescent matorral**, **T24 Olea europaea-Ceratonia siliqua forest**, **T3A Mediterranean lowland to submontane Pinus forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz