

Plantago sempervirens

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **10.27**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.36**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7.3**

Moisture indicator value: **2**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.18**
Disturbance frequency (herb layer): **1.7**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.46**
Grazing pressure: **0.3**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Constant species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**