

# ***Plantago schwarzenbergiana***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **8**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **7**

### Disturbance indicator values

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **2.09**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.26**

Grazing pressure: **0.53**

Soil disturbance: **0.23**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DB \(FEP\) Festuco-Puccinellietea](#)

### Broad habitat

Occurrence in broad habitats: **Saline vegetation**