

Pleurospermum austriacum

Habitus and growth type

Plant height [m]: **0.99**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **4.78**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.1**

Temperature indicator value: **3.7**

Moisture indicator value: **5.9**

Reaction indicator value: **7**

Nutrient indicator value: **5.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.26**

Disturbance frequency (herb layer): **1.28**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.07**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#), [DA \(FES\) *Festuco-Brometea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **6**

Continentality amplitude: **6**