Poa timoleontis

Habitus and growth type

Plant height [m]: **0.16**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **8.1**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.92**
Disturbance frequency (herb layer): **1.7**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.14**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.35**
Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FR (DAP) *Daphno-Festucetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S75 Eastern Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz