

# *Poa timoleontis*

## Habitus and growth type

Plant height [m]: **0.16**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **8.1**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.14**

Grazing pressure: **0.35**

Soil disturbance: **0.24**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*FR \(DAP\) \*Daphno-Festucetea\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),  
Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**