

Potentilla anglica

Habitus and growth type

Plant height [m]: **0.27**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **25.79**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.57**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **5.8**

Reaction indicator value: **5.8**

Nutrient indicator value: **3.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.29**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.71**

Grazing pressure: **0.32**

Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **3**

Continentality amplitude: **3**