

# *Potentilla brauneana*

## Habitus and growth type

Plant height [m]: **0.04**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **1.5**

Moisture indicator value: **7**

Reaction indicator value: **9**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.34**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.23**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.03**

Grazing pressure: **0.21**

Soil disturbance: **0.16**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R41 Snow-bed vegetation](#)

Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **2**