

# *Potentilla nitida*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **2**

Moisture indicator value: **2**

Reaction indicator value: **9**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **0.77**

Disturbance severity: **0.21**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0**  
Grazing pressure: **0.14**  
Soil disturbance: **0.05**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [U36 Temperate high-mountain base-rich inland cliff](#)

Constant species of EUNIS habitats: [U36 Temperate high-mountain base-rich inland cliff](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**