

Potentilla reptans

Habitus and growth type

Plant height [m]: **0.23**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **24.54**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.6**

Reaction indicator value: **6.9**

Nutrient indicator value: **5.2**

Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.31**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.8**

Grazing pressure: **0.26**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R34 Submediterranean moist meadow](#), [R36 Moist or wet mesotrophic to eutrophic pasture](#)

Constant species of EUNIS habitats: [R31 Mediterranean tall humid inland grassland](#), [R34 Submediterranean moist meadow](#), [R36 Moist or wet mesotrophic to eutrophic pasture](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Scrub, Synanthropic**

Distribution

Continentality amplitude: **6**