

# *Primula deorum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Moisture indicator value: **7**

Disturbance indicator values

Disturbance frequency: **1.18**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.13**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.11**

Grazing pressure: **0.11**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*Q23 Relict mire of Mediterranean mountains\*\*](#)

Constant species of EUNIS habitats: [\*\*Q23 Relict mire of Mediterranean mountains\*\*](#)

Broad habitat

Occurrence in broad habitats: **Mire**