

# **Psephellus sibiricus**

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Disturbance indicator values

Disturbance frequency: **1.63**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.87**

Grazing pressure: **0.42**

Soil disturbance: **0.17**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*BC \(BRA\) Brachypodio pinnatifidae\*\*](#), [\*\*Betuletea pendulae\*\*](#), [\*\*DA \(FES\) Festuco-Brometea\*\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*U35 Boreal and Arctic base-rich inland cliff\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**