Pseudotsuga menziesii

Habitus and growth type

Plant height [m]: **45.09**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Leaf

Leaf type (woody plants only): **Needle-like**
Leaf phenology (woody plants only): **Evergreen**
Specific leaf area [mm^2/mg]: **7.17**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **426.69**
Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **4, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Disturbance indicator values

Disturbance frequency: **0.08**
Disturbance frequency (herb layer): **1.3**
Disturbance severity: **0.83**
Disturbance severity (herb layer): **0.18**
Mowing frequency: **0.01**
Grazing pressure: **0.2**
Soil disturbance: **0.24**

Habitat and sociology

EUNIS habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of EUNIS habitats: **T3M Coniferous plantation of non site-native trees**

Constant species of EUNIS habitats: **T3M Coniferous plantation of non site-native trees**

Dominant species of EUNIS habitats: **T3M Coniferous plantation of non site-native trees**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz