

# *Ptilostemon chamaepeuce*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

## Fruit, seed and dispersal

Seed mass [mg]: **11.81**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **11**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.48**

Disturbance frequency (herb layer): **1.45**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.29**

Grazing pressure: **0.38**

Soil disturbance: **0.17**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**