

# ***Pulicaria odora***

## **Habitus and growth type**

Plant height [m]: **0.58**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **May-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.1**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7**

Temperature indicator value: **8.8**

Moisture indicator value: **4.5**

Reaction indicator value: **6**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.19**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.24**

Mowing frequency: **0.03**

Grazing pressure: **0.27**

Soil disturbance: **0.1**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

### EUNIS habitat

Constant species of EUNIS habitats: [S51 Mediterranean maquis and arborescent matorral](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**