

Pulmonaria angustifolia

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **17.27**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.5**

Temperature indicator value: **6.5**

Moisture indicator value: **4.9**

Reaction indicator value: **6.3**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**

Disturbance frequency (herb layer): **1.55**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.26**

Grazing pressure: **0.24**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) *Quercetea pubescentis*](#), [CL \(GER\) *Trifolio-Geranietea sanguinei*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**

Distribution

Continentality: **6**

Continentality amplitude: **3**