

Pyrola minor

Habitus and growth type

Plant height [m]: **0.16**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **21.91**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **3.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.15**

Disturbance frequency (herb layer): **1.13**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BA \(PIC\) Vaccinio-Piceetea](#), [CA \(FAG\) Carpino-Fagetea sylvaticae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [U21 Boreal and Arctic siliceous scree and block field](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [U21 Boreal and Arctic siliceous scree and block field](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **5**