Pyrola rotundifolia

Habitus and growth type

Plant height [m]: **0.19**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.95**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.28**
Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.3**
Temperature indicator value: **4**
Moisture indicator value: **6.1**
Reaction indicator value: **5.4**
Nutrient indicator value: **3.1**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.18**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **0.99**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0.02**
Grazing pressure: **0.2**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BA (PIC) *Vaccinio-Piceetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3H Larix light taiga**, **U35 Boreal and Arctic base-rich inland cliff**

Constant species of EUNIS habitats: **T3H Larix light taiga**, **U35 Boreal and Arctic base-rich inland cliff**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz