

Quercus rotundifolia

Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **5.15**

Fruit, seed and dispersal

Dispersal mode: **Dyszoochory, Anthropochory**

Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8**

Moisture indicator value: **3**

Reaction indicator value: **6.5**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.09**

Disturbance frequency (herb layer): **1.3**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.14**

Mowing frequency: **0.01**

Grazing pressure: **0.28**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T21 Mediterranean evergreen Quercus forest](#)

Constant species of EUNIS habitats: [S62 Western acidophilous garrigue](#), [T21 Mediterranean evergreen Quercus forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**