Quercus suber

Habitus and growth type

Plant height [m]: **15.78**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **6.12**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **3504.4**
Dispersal mode: **Dyszoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **8**
Moisture indicator value: **3**
Reaction indicator value: **3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.06**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0**
Grazing pressure: **0.27**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**, **T21 Mediterranean evergreen Quercus forest**
Constant species of EUNIS habitats: **T21 Mediterranean evergreen Quercus forest**
Dominant species of EUNIS habitats: **T21 Mediterranean evergreen Quercus forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz