

Quercus suber

Habitus and growth type

Plant height [m]: **15.78**

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **6.12**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **3504.4**

Dispersal mode: **Dyszoochory, Anthropochory**

Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **8**

Moisture indicator value: **3**

Reaction indicator value: **3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0**
Grazing pressure: **0.27**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S62 Western acidophilous garrigue](#), [T21 Mediterranean evergreen Quercus forest](#)

Constant species of EUNIS habitats: [T21 Mediterranean evergreen Quercus forest](#)

Dominant species of EUNIS habitats: [T21 Mediterranean evergreen Quercus forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**