

Ranunculus aconitifolius

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **36.19**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.56**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.6**

Temperature indicator value: **3.7**

Moisture indicator value: **7.5**

Reaction indicator value: **5.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.13**

Grazing pressure: **0.18**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R56 Montane to subalpine moist or wet tall-herb and fern fringe](#)

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Forest**

Distribution

Continentality: **4**

Continentality amplitude: **2**