

# ***Ranunculus marschlinii***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **3**

Moisture indicator value: **7**

Reaction indicator value: **7**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.63**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0**

Grazing pressure: **0.2**

Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [OA \(MON\) Montio-Cardaminetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [U24 Mediterranean siliceous scree](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Sparsely vegetated (incl. rock and scree)**