Ranunculus oxyspermus

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.59**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.3**
Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **1.87**
Disturbance frequency (herb layer): **2.26**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.38**
Grazing pressure: **0.42**
Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1C Desert steppe**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz